



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/18, 15/19, 15/24, 15/25, A61K 38/18, 38/19, 38/20		(11) International Publication Number: WO 95/08630
A1		(43) International Publication Date: 30 March 1995 (30.03.95)
(21) International Application Number: PCT/US94/10800		(81) Designated States: AU, CA, JP, KR, NZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
(22) International Filing Date: 23 September 1994 (23.09.94)		
(30) Priority Data: 08/126,973 24 September 1993 (24.09.93) US 08/290,373 15 August 1994 (15.08.94) US		Published With international search report.
(71) Applicants: AMERICAN CYANAMID COMPANY [US/US]; One Cyanamid Plaza, Wayne, NJ 07470-8426 (US). YEDA RESEARCH AND DEVELOPMENT CO., LTD. [IL/IL]; The Weizmann Institute of Science, P.O. Box 95, 76100 Rehovot (IL).		
(72) Inventors: SEDDON, Andrew, Peter; 71 Summit Street, Monroe, NY (US). LI, Lu-Yuan; 20 Gerardine Place, New City, NY 10956 (US). BÖHLEN, Peter; Two Amalfi Drive, Peekskill, NY (US). EISINGER, Magdalena; 30 Pine Terrace, Demarest, NJ (US). YAYON, Avner; Weizmann Institute of Science, 76100 Rehovot (IL).		
(74) Agents: KRINSKY, Mary, M. et al.; St. Onge Steward Johnston & Reens, 986 Bedford Street, Stamford, CT 06905 (US).		
(54) Title: SURFACE LOOP STRUCTURAL ANALOGUES OF FIBROBLAST GROWTH FACTORS		
<p>FGF-2 Asn Asn Tyr Asn Thr Tyr Arg Ser Arg Lys Tyr _____ Thr Ser Trp Tyr Val Ala Leu Lys 110 115 120 (Ser) 125</p> <p>FGF-1 Asn His Tyr Asn Thr Tyr Ile Ser Lys Lys His Ala Glu Lys Asn Trp Phe Val Gly Leu Lys (His)</p> <p>IL-18 Asn Asn Lys Leu Glu Phe Glu Ser Ala Glu Phe _____ Pro Asn Trp Tyr Ile Ser Thr Ser</p>		
(57) Abstract		
<p>Structural analogues of fibroblast growth factors have an amino acid sequence replacement in the ninth or tenth beta strand of the factor, or the sequence that corresponds to the surface loop that connects the ninth and tenth beta strands, such that the folding of the molecule is not significantly perturbed. Preferred analogues have the overall secondary and tertiary structure of the original factor, and bind to heparin and a member or members of the fibroblast growth factor receptor family with high affinity. In some embodiments the replacement is with another amino acid sequence such as a loop sequence from another structurally related fibroblast growth factor or an interleukin. The figure discloses amino acid sequences from FGF-1, FGF-2, and IL-1 beta.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

SURFACE LOOP STRUCTURAL ANALOGUES
OF FIBROBLAST GROWTH FACTORS

Related U.S. Application Data

5 This is a continuation-in-part of co-pending U.S. patent application serial number 08/126,973, filed September 24, 1993.

Technical Field of the Invention

10 This invention relates to fibroblast growth factor analogues having different primary structures in a surface loop that connects the ninth and tenth β -strands. The analogues retain overall secondary and tertiary protein structural similarities to the original factors but exhibit different biological properties and receptor
15 binding specificity profiles.

Background of the Invention

 Polypeptide growth factors are hormone-like modulators of cell proliferation and differentiation. Growth factors are responsible for the regulation of a variety
20 of physiological processes, including development, regeneration, and wound repair, and have been associated with normal as well as with pathophysiological processes. Numerous growth factors have been identified in various tissues and cells, and names that have been applied to
25 these factors include epidermal growth factor, platelet-derived growth factor, nerve growth factor, hematopoietic growth factors, and fibroblast growth factor.

- 2 -

Fibroblast growth factor (FGF) was first described as an activity derived from bovine brain or pituitary tissue which was mitogenic for fibroblasts and endothelial cells. It was later noted that the primary mitogen
5 from brain was different from that isolated from pituitary. These two factors were named acidic and basic FGF, respectively, because they had similar biological activities but differed in their isoelectric points. Acidic and basic FGF are proteins containing approximately
10 ly 154 amino acids. Their amino acid sequences are related, with approximately 55% sequence identity between them.

Acidic and basic fibroblast growth factors are now known to be members of a larger family of heparin-binding
15 growth factors that collectively trigger a variety of biological responses in many cell types, including those of mesoderm and neuroectoderm origin, such as endothelial cells, smooth muscle cells, adrenal cortex cells, prostatic and retina epithelial cells, oligodendrocytes,
20 astrocytes, chondrocytes, myoblasts, and osteoblasts. As original family members, acidic and basic FGF are now denoted FGF-1 and FGF-2, respectively. Seven other members of the family have been identified on the basis of their modulation of cell proliferation and differentiation,
25 and their sequence homology to other FGFs.

In addition to eliciting a mitogenic response that stimulates cell growth, fibroblast growth factors can stimulate a large number of cell types to respond in a non-mitogenic manner. These activities include promotion
30 of cell migration into wound areas (chemotaxis), initiation of new blood vessel formulation (angiogenesis), modulation of nerve regeneration and survival (neurotrophism), modulation of endocrine functions, and stimulation or suppression of specific cellular protein expression,
35 extracellular matrix production and cell survival

- 3 -

(Baird, A., and Böhlen, P., *Handbook of Exp. Pharmacol.* 95(1): 369-418, Springer, 1990). These properties provide a basis for using fibroblast growth factors in therapeutic approaches to accelerate wound healing, nerve repair, collateral blood vessel formation, and the like. For example, fibroblast growth factors have been suggested to minimize myocardium damage in heart disease and surgery (U.S. Pat. No. 4,378,347 to Franco).

Current research regarding FGF-2 and other FGFs has centered on the molecular details of the receptor-mediated pathways by which their diverse physiological activities are expressed, as a way to gain information for the design of therapeutically useful agents that can either mimic or inhibit the action of these factors. Since the primary structure of FGF-2 isolated from a variety of sources is known, and bovine and human FGF-2 have been cloned and expressed in *E. coli* and *S. cerevisiae*, recent attention has focused on secondary and tertiary structure.

The 3-dimensional structures of FGF-1 and FGF-2 have been determined (Eriksson, E.A., et al., *Proc. Nat. Acad. Sci. USA* 88: 3441-3445 (1991), Zhang, J., et al., *Proc. Nat. Acad. Sci. USA* 88: 3446-3450 (1991), and Zhu, H., et al., *Science* 251: 90-93 (1991)). In these studies, FGF-1 and FGF-2 were shown to exhibit a folding pattern strikingly similar to that observed for the cytokine interleukin-1 α , and interleukin-1 β (IL-1 α and IL-1 β), protein factors produced by macrophages and T-cells in response to antigenic or mitogenic stimulation, though the primary structures of interleukin-1 polypeptides have only about a 10% amino acid sequence correspondence to FGFs.

The overall structure of FGF-2 can be described as a trigonal pyramid where each of the three sides are built of two β -strands together forming a β -sheet barrel of six

- 4 -

antiparallel strands (Eriksson, E.A., et al., *Proc. Nat. Acad. Sci. USA* 88: 3441-3445 (1991)). The base of the pyramid is built of six additional β -strands extending from the three sides of the pyramid to close one end of the barrel for a total of twelve β -strands. Thus, a threefold repeat is observed in the folding of the polypeptide chain and a pseudo-three-fold axis passes through the center of the base of the molecule and extends through the apex of the pyramid (*ibid.*). Of the amino acids conserved within the FGF family of proteins, most are located within the core β -strand regions of FGF-2, supporting the expectation that each of these proteins has an overall 3-dimensional structure similar to that of FGF-2.

The biological responses of FGF are mediated by the heparan sulfate-dependent binding of the growth factor to specific cell surface receptors (Givol, D., and Yayon, A., *FASEB J.* 6: 3362-3369 (1992) and Jaye, M., et al., *Biochim. Biophys. Acta* 1135: 185-199 (1992)), yet the molecular interactions of heparin and receptor with FGF and the exact nature of the events of the signal transduction pathway are unknown. Studies employing synthetic peptides related to the FGF sequence showed that FGF-2 (33-77) and (106-129) bind to heparin and act as weak partial agonists and antagonists in a mitogenic assay of FGF activity (Baird, A., et al. *Proc. Nat. Acad. Sci. USA* 85: 2324-2328 (1988)). The same study identified a sequence, FGF-2 (115-124), involved in receptor binding. The segment begins in the middle of the ninth β -strand, makes a somewhat open loop on the surface of the folded molecule, and terminates at the beginning of the tenth β -strand. This sequence (118-122) in the native protein forms a small surface loop that is close to a cluster of basic surface residues that may form a putative heparin binding site (Zhang, J., et al., *Proc. Nat. Acad. Sci. USA* 88: 3446-3450 (1991)).

- 5 -

This sequence also contains Thr-121, which can be phosphorylated by a cAMP-dependent protein kinase (Feige, J.J., and Baird, A., *Proc. Nat. Acad. Sci. USA* 86: 3174-3178 (1989)); Thr-121 is denoted in the paper as Thr-112 since the investigators employed the N-terminally truncated form of the polypeptide, which exhibits full biologic activity but has 146 rather than the usual 154 amino acids). Phosphorylation of Thr-121 results in the generation of a form of the protein that exhibits an increased capacity to compete with radiolabelled FGF-2 binding to its receptor, but no difference in the biological properties of the phosphorylated and nonphosphorylated forms of the protein were observed (*ibid.*). In contrast to FGF-2, FGF-1 is not a substrate for the kinase. The data are consistent with the hypothesis that the sequence 115-124 is involved in receptor binding, but they do not define the complete receptor binding domain of the molecule, nor do they demonstrate the physiological significance of phosphorylation of Thr-121.

In addition to heparin and receptor binding regions, there is evidence that specific sequences in FGF influence ligand-induced signal transduction. For example, deletion of residues 27-32 of FGF-2 (Lys-Asp-Pro-Arg-Leu) or mutation of the basic residues Arg-118, Lys-119, Lys-128, and Arg-129 did not appear to affect the mitogenic activity of the protein, but eliminated activation of plasminogen activator gene expression (Eur. Pat. Ap. Pub. No. 363,675 to Bergonzoni, L., et al., and Presta, M., et al., *Biochem. Biophys. Res. Com.* 185: 1098-1107 (1992)).

At least four different fibroblast growth factor receptors (FGFR) have been identified (Werner, S., et al., *Mol. Cell. Bio.* 12: 82-88 (1992)), and functional differences between different receptor forms have been observed. Different FGF receptor forms derived from the same gene via alternative splicing have different ligand

- 6 -

binding properties, and analogous splice variants from different FGF receptor genes bind different members of the FGF family (Johnson, D.E., and Williams, L.T., *Adv. Can. Res.* 60: 1-41 (1993)). Thus, fibroblast growth factor receptors exhibit a multitude of structural variants, and considerable cross-reactivity between receptors and their various ligands (Yayon, A., et al., *EMBO J.* 11: 1885-1890 (1992)). Though the carboxy-terminal region of the third immunoglobulin-like domain appears to be a structural element that defines specificity of different FGF members (Werner, et al., and Yayon, et al., cited above), the precise nature of FGF-receptor-heparin interactions and the protein residues involved have yet to be elucidated.

15

Summary of the Invention

It is an objective of the invention to provide new structural analogues of fibroblast growth factors.

It is another objective of the invention to provide fibroblast growth factor analogues having binding and biological activities that are pharmacologically dissociated, i.e., that bind to an FGF receptor, which may be the same or different from a receptor that normally binds the wild-type factor, and/or exhibits different biological properties from that observed in the wild-type factor.

It is a further and more specific objective of the invention to provide fibroblast growth factor structural analogues such as structural analogues of human fibroblast growth factor-2, which have all or part of a surface loop replaced with another amino acid sequence.

These and other objectives are accomplished by the present invention which provides fibroblast growth factor

- 7 -

structural analogues having amino acid changes in surface loops, or sequences adjacent to the surface loops, such that the overall secondary and tertiary structures of the polypeptides are not significantly perturbed, and the analogues bind to heparin. More particularly, this invention provides fibroblast growth factor structural analogues having an amino acid sequence replacement in the ninth or tenth β -strand of the factor, or the sequence that corresponds to the surface loop that connects the ninth and tenth β -strands, such that the interaction of β -strands nine and ten with adjacent strands is not significantly perturbed, and the analogue binds to heparin. The analogues exhibit binding affinity to a fibroblast growth factor receptor. Binding specificity of the analogues is preferably different from the binding specificity of the native factor to the corresponding receptor. In one embodiment, the analogue stimulates *in vivo* angiogenesis.

Preferred fibroblast growth factor structural analogues have an amino acid sequence substitution in the surface loop that extends from the ninth to the tenth β -strand. In some embodiments, all or part of the amino acid sequence in the growth factor surface loop extending from the ninth to the tenth β -strands is replaced by another amino acid sequence such as a corresponding amino acid sequence derived from related fibroblast growth factor or interleukin polypeptides.

Thus, the fibroblast growth factor analogues of this invention include, but are not limited to, analogues formed by using, in replacement of corresponding loop sequences in other growth factors, fibroblast growth factor-1 sequence Lys115-Lys116-His117-Ala118-Glu119-Lys120-Asn121 (amino acids 115-121 of SEQ ID NO 2), fibroblast growth factor-2 sequence Arg118-Lys119-Tyr120-Thr121-Ser122 (amino acids 118-122 of SEQ ID NO 1), fi-

- 8 -

broblast growth factor-3 sequence Arg-Leu-Tyr-Arg-Thr-Val-Ser-Ser-Thr-Pro-Gly-Ala-Arg-Arg-Gln-Pro-Ser-Ala-Glu-Arg-Leu (amino acids 132-152 of SEQ ID NO 4), fibroblast growth factor-4 sequence Tyr-Lys-Tyr-Pro-Gly (amino acids 172-176 of SEQ ID NO 5), fibroblast growth factor-5 sequence Ala-Ile-His-Arg-Thr-Glu-Lys-Thr-Gly-Arg-Glu (amino acids 176-186 of SEQ ID NO 6), fibroblast growth factor-6 sequence Asp-Leu-Tyr-Gln-Gly (amino acids 174-178 of SEQ ID NO 7), fibroblast growth factor-7 sequence Ala-Lys-Trp-Thr-His-Asn-Gly-Gly-Glu (amino acids 154-163 of SEQ ID NO 8), fibroblast growth factor-8 sequence Ala-Lys-Tyr-Glu-Gly (amino acids 144-148 of SEQ ID NO 9), and fibroblast growth factor-9 sequence Asn-Leu-Tyr-Lys-His-Val-Asp-Thr-Gly-Arg-Arg (amino acids 151-161 of SEQ ID NO 10). Analogues having loop flanking sequences changed, and peptide fragment analogues containing the loop sequences are correspondingly provided by the invention.

For example, human fibroblast growth factor-2 structural analogues having amino acid sequence substitutions in residues 112 to 128 result in factors that retain the same overall secondary and tertiary structure as FGF-2, but exhibit different biological properties. In these exemplary embodiments, an FGF-2 loop sequence is replaced with any size or composition of insert. Some derivatives are prepared by changing residues 115 to 124; others are prepared by changing loop residues 118 to 122. In one embodiment, for example, the FGF-2 loop is replaced with corresponding amino acids derived from FGF-1 from the same or another species; in another, an interleukin-1 β from the same or another species is used; and, in another, a loop from FGF-7 from the same or another species is used. As summarized above, these analogues bind heparin and a native fibroblast growth factor receptor.

The invention also provides DNA encoding the fibroblast growth factor derivatives, biologically functional

- 9 -

circular plasmid or viral DNA vectors comprising the DNA, and procaryotic or eucaryotic host cells such as *E. coli* transformed or transfected with the vectors in a manner allowing the host cell to express the new factors.

5 Brief Description of the Figures

Figure 1 sets out sequence alignment of human FGF-2, FGF-1 and IL-1 β in the loop region (identified in the Sequence Listing section hereinafter as ID NOs 1 to 3).

10 The sequences underlined correspond to the surface loop located at the end of the ninth β -strand. The residues in parenthesis indicate the residues found in the bovine sequences.

Figure 2 plots competitive receptor binding of confluent baby hamster kidney cells incubated with ¹²⁵I-FGF-2 and indicated concentrations of competitors FGF-2 (open circles), structural analogue FGF-2LI (described in greater detail hereinafter, closed squares), and structural analogue FGF-2LA (described hereinafter, closed triangles).

20 Figure 3 plots the binding of radioiodinated FGF-2 (closed squares), FGF-2LA (open circles), and FGF-2LI (open triangles) to FGF-receptor type 1 present on NIH 3T3 cells as a function of added unlabelled FGF proteins.

Figure 4 shows endothelial cell growth of adult bovine aortic arch endothelial cells plated at 8000 cells/well and incubated for 5 days with the indicated concentrations of FGF-2 (closed circles), FGF-2LI (open squares), and FGF-2LA (open triangles).

Figure 5 is a bar graph depicting production of cell-associated urokinase-type plasminogen activator induced by wild-type or mutant FGF-2 (10 ng/ml).

30

- 10 -

Figure 6 is a bar graph illustrating the potency of *in vitro* angiogenic activities of wild-type or mutant FGF-2 (10 ng/ml) as measured by their abilities to induce formation of tube-like structures by ABAE cells cultured on a type I collagen gel. The data are from a typical experiment in which cell cultures of identical conditions are maintained in duplicate wells. Three to four areas of each well are examined by image analysis and the mean value and standard deviation are presented.

Figure 7 presents an amino acid sequence comparison among the nine members of the FGF family between residues corresponding to amino acid residues 113 and 128 in FGF-2 (SEQ ID NO 1; FGF-1 is set out as SEQ ID NO 2, and the other FGFs are sequentially set out as SEQ ID NOS 4 to 10, except that His121 of the FGF-1 bovine sequence is replaced by Asn in the human sequence). The numbering system is relative to FGF-2 and the asterisks refer to identical and conserved residues. Residues located between the FGF-2 locations Ser117 to Trp123 mark the positions where other FGFs contain inserts and changes. To illustrate homologous sequences in the ninth and tenth β -strands and in the loop region, the figure employs standard one-letter nomenclature for the amino acids: A, Ala; C, Cys; D, Asp; E, Glu; F, Phe; G, Gly; H, His; I, Ile; K, Lys; L, Leu; M, Met; N, Asn; P, Pro; Q, Gln; R, Arg; S, Ser; T, Thr; V, Val; W, Trp; and Y, Tyr.

Figure 8 shows a comparison of the degree of neo-vascularization induced in a rabbit ear chamber model by FGF-2 (closed triangles) and FGF-2LA (closed circles). A control without FGF is represented by closed squares.

Figure 9A plots the binding of ^{125}I -FGF-2 as a function of added FGF-1 (open circles), FGF-2 (open squares), FGF-7 (closed circles), FGF-2LA (closed squares), and FGF-2LI (open triangles) to soluble recombinant FGF-re-

- 11 -

ceptor type 1 as a function of added unlabelled FGF proteins. Figure 9B plots a similar binding profile, except that soluble recombinant FGF-2(IIIc) is employed.

Detailed Description of the Invention

5 This invention is based upon the finding that changes in fibroblast growth factor surface loop residues in a manner that does not perturb the overall secondary and tertiary structure of the FGF molecule yields an array of structural analogues having varied and desirable physiological properties, including FGF agonists and potential
10 antagonists. Changes in loop sequences, or sequences adjacent to loop sequences, yield analogues that bind to heparin and exhibit binding affinity to a native fibroblast growth factor receptor.

15 By "antagonist" is meant any substance that tends to nullify the action of another, as one, for example, that binds to a cell receptor without eliciting a biological response. For FGFs, antagonists include, but are not limited to, any substance that binds to an FGF receptor
20 but does not stimulate proliferation or migration of fibroblasts, endothelial cells and astroglial cells. In contrast, by "agonist" is meant any substance that has affinity for and stimulates physiological activity at cell receptors normally stimulated by naturally occurring
25 factors, including mitogenic activity, angiogenesis, chemotaxis, and the like. As used herein, the terms are not limiting, and a single factor may exhibit antagonism to one biological function mediated by native factor, and agonism to another function. Where this occurs, the
30 biological activities of native factor are said to be pharmacologically dissociated in the analogue.

A protein is defined herein as a fibroblast growth factor (FGF) if it shows significant sequence and three-

- 12 -

dimensional structural homology to other members of the FGF family, FGF-like activity in *in vitro* or *in vivo* assays and binds to heparin or heparin-like substances. By "heparin" is meant the heterogeneous, sulfated anionic polysaccharide composed of D-glucuronic acid and D-glucosamine, bound to a protein core as the "proteoglycan" or in a free form as the "aglycan", that may or may not have anticoagulant properties. "Heparin-like" substances are molecules that have oligosaccharide structures related to the heparins, but may or may not have anticoagulant activity. Any type of fibroblast growth factor or mutein or derivative is encompassed by this invention, particularly human fibroblast growth factor, as well as loop sequence peptides that exhibit antagonist or agonist properties, or both.

Broadly speaking, the fibroblast growth factor analogues of this invention include structural analogues of fibroblast growth factor having at least one amino acid deletion, insertion or replacement in a β -strand, or a loop connecting two β -strands such that adjacent strands are not perturbed, and the molecule retains its overall secondary and tertiary structure. Secondary and tertiary structure is determined by heparin binding, binding to an FGF receptor, spectroscopy (using ultraviolet, visible, or circular dichroic light), X-ray crystallography, and biological activities.

Particularly preferred are FGF sequence replacements in a loop connecting two β -strands, analogous to the structural cassette or module replacements of turns with retention of parent conformation disclosed for enzymes by Hynes, T.R., et al., *Nature* 339: 73-76 (1989). In some embodiments, the ninth or tenth β -strand of the factor, or the sequence that corresponds to the surface loop that connects the ninth and tenth β -strands, are replaced with any size or composition amino acid sequence insert such

- 13 -

that overall molecular folding is not significantly perturbed. In certain embodiments, the surface loop is replaced with a surface loop from another factor. Thus, this invention specifically encompasses loop and loop stem and anchor replacements with structurally related and unrelated sequences, including random loops, longer or shorter loops, and differently charged loops and their stems and anchors. It also encompasses peptide fragments that mimic these loops in binding studies and biological assays.

As set out more fully below, numbering of the amino acids in these strands is, for convenience, made with reference to FGF-2. Perturbation and strand adjacency are herein defined in a three-dimensional sense and measured as described above. Overall folding is maintained in the analogues. Preferred analogues bind to an FGF receptor with an affinity that is greater than, or up to 100-times lower than the binding of a corresponding native FGF family member exhibiting affinity to the receptor.

Receptors that many of the analogues typically bind include native baby hamster kidney (BHK) cell fibroblast growth factor receptor, NIH 3T3 cell receptors, or other receptors specific to particular FGFs. Binding preferably exhibits an affinity substantially similar or superior to the binding of the corresponding native factor to that receptor. An advantage of the invention is that the loop manipulation results in analogues exhibiting similar binding profiles to the corresponding factors from which the loops are derived. Thus, certain analogues of the invention target receptors that the native factor does not bind, including receptors on different cell types. For example, factors that ordinarily bind to endothelial cell receptors can be engineered to bind to epithelial cell receptors, and so forth. Examples are given herein-

- 14 -

after. One embodiment stimulates angiogenic activity *in vivo*.

Especially preferred structural analogues are those in which the factor has at least one amino acid deletion, addition or substitution in the sequence of the surface loop that extends from the ninth to the tenth β -strand and protrudes from the surface when the molecule is folded as described by Eriksson, et al., Zhu, et al., and Zhang, et al., cited above, and has the overall secondary and tertiary structure of the native factor. In some embodiments, the replacements are sequences from other FGF species or related molecules. In other embodiments, the replacements are structurally unrelated to any FGF species, and may contain fewer or more amino acids, or differently charged amino acids. Thus, any amino acid sequence can be used as a sequence replacement in the loop connecting the strands.

For example, structural FGF analogues of the invention include analogues having at least one amino acid deletion, addition or substitution, particularly amino acid sequence substitution, in surface loops that include, but are not limited to, fibroblast growth factor-1 sequence 112 to 123, particularly Lys115-Lys116-His117-Ala118-Glu119-Lys120-Asn121 (amino acids 115-121 of SEQ ID NO 2); fibroblast growth factor-2 sequence 115 to 124, particularly Arg118-Lys119-Tyr120-Thr121-Ser122 (amino acids 115-124 of SEQ ID NO 1), fibroblast growth factor-3 sequence Arg-Leu-Tyr-Arg-Thr-Val-Ser-Ser-Thr-Pro-Gly-Ala-Arg-Arg-Gln-Pro-Ser-Ala-Glu-Arg-Leu (amino acids 132-152 of SEQ ID NO 4), fibroblast growth factor-4 sequence Tyr-Lys-Tyr-Pro-Gly (amino acids 172-176 of SEQ ID NO 5), fibroblast growth factor-5 sequence Ala-Ile-His-Arg-Thr-Glu-Lys-Thr-Gly-Arg-Glu (amino acids 176-186 of SEQ ID NO 6), fibroblast growth factor-6 sequence Asp-Leu-Tyr-Gln-Gly (amino acids 174-178 of SEQ ID NO 7),

- 15 -

fibroblast growth factor-7 (also known as keratinocyte growth factor or KGF) sequence Ala-Lys-Trp-Thr-His-Asn-Gly-Gly-Glu (amino acids 154-162 of SEQ ID NO 8), fibroblast growth factor-8 sequence Ala-Lys-Tyr-Glu-Gly (amino acids 144-148 of SEQ ID NO 9), and fibroblast growth factor-9 sequence Asn-Leu-Tyr-Lys-His-Val-Asp-Thr-Gly-Arg-Arg (amino acids 151-161 of SEQ ID NO 10). A comparison of surface loop sequences in the FGF family is depicted in Figure 7. In these embodiments of the invention, structural analogues are prepared by replacing the surface loop sequence of one FGF with another amino acid sequence such as, but not limited to, a surface loop sequence from another factor.

Changes in neighboring regions that affect the loop, including loop stems and anchor points, are encompassed by the invention, so long as the overall folding of the molecule is not perturbed. Thus, analogues of the invention include structures having mutations, particularly sequence replacements, in fibroblast growth factor-1 sequence Tyr112-Ile113-Ser114-Lys115-Lys116-His117-Ala118-Glu119-Lys120-Asn121-Trp122-Phe123-Lys124-Asn125 (amino acids 112-125 of SEQ ID NO 2), fibroblast growth factor-2 sequence Tyr115-Arg116-Ser117-Arg118-Lys119-Tyr120-Thr121-Ser122-Trp123-Tyr124 (amino acids 115-124 of SEQ ID NO 1), fibroblast growth factor-3 sequence Tyr-Ala-Ser-Arg-Leu-Tyr-Arg-Thr-Val-Ser-Ser-Thr-Pro-Gly-Ala-Arg-Arg-Gln-Pro-Ser-Ala-Glu-Arg-Leu-Trp-Tyr (amino acids 129-154 of SEQ ID NO 4), fibroblast growth factor-4 sequence Tyr-Glu-Ser-Tyr-Lys-Tyr-Pro-Gly-Met-Phe (amino acids 169-178 of SEQ ID NO 5), fibroblast growth factor-5 sequence Tyr-Ala-Ser-Ala-Ile-His-Arg-Thr-Glu-Lys-Thr-Gly-Arg-Glu-Trp-Tyr (amino acids 173-188 of SEQ ID NO 6), fibroblast growth factor-6 sequence Tyr-Glu-Ser-Asp-Leu-Tyr-Gln-Gly-Thr-Tyr (amino acids 171-180 of SEQ ID NO 7), fibroblast growth factor-7 sequence Tyr-Ala-Ser-Ala-Lys-Trp-Thr-His-Asn-Gly-Gly-Glu-Met-Phe (amino acids 151-164

- 16 -

of SEQ ID NO 8), fibroblast growth factor-8 sequence Asn-Asn-Tyr-Thr-Ala-Leu-Gln-Asn-Ala-Lys-Tyr-Glu-Gly-Trp-Tyr-Met-Ala-Phe-Thr-Arg-Lys (SEQ ID NO 9), and fibroblast growth factor-9 sequence Asn-Trp-Tyr-Asn-Thr-Tyr-Ser-Ser-Asn-Leu-Tyr-Lys-His-Val-Asp-Thr-Gly-Arg-Arg-Tyr-Tyr-Val-Ala-Leu-Asn-Lys-Asp (SEQ ID NO 10). The invention also encompasses peptides corresponding to these loops that exhibit heparin and FGF receptor binding.

Structural analogues of FGF-2 are preferred in some embodiments. By "FGF-2" is meant any fibroblast growth factor-2 exhibiting biologic activity including the 146-amino acid polypeptide originally isolated and sequenced, the 154 amino acid form currently thought to be the full polypeptide, truncated forms exhibiting activity, extended forms such as placental FGF, higher molecular weight N-terminally extended forms described in the literature and analogues including derivatives and muteins of any of these. The term specifically includes natural FGF-2 extracted from mammalian tissue as well as recombinant polypeptides expressed from cloned DNA in *E. coli* or *S. cerevisiae* from any species or expressed in insect or mammalian cells with appropriate vectors.

Human FGF-2 is preferred in many embodiments. Human FGF-2 having 9th or 10th β -strands which can be manipulated according to the invention includes, but is not limited to, FGF-2 having amino acid additions, amino acid substitutions, and amino acid deletions, including deletions of portions of the amino or carboxyl terminus, and chimeric proteins containing FGF at the N-or C-terminus of another protein. Example FGF-2s which can be manipulated according to the invention include those having cysteine substituted with a neutral amino acid such as serine, or aspartic acid, arginine, glycine, serine, or valine substituted with other acids suggested to have enhanced stability in Eur. Pat. Ap. Pub. No. 281,822 to

- 17 -

Seno, et al.; muteins formed by replacing at least one, and more preferably two, of the cysteines found in natural FGF-2 with a different amino acid residue to yield a more stable analogue (Eur. Pat. Ap. Pub. No. 320,148 to Arakawa and Fox); muteins lacking amino acids from the carboxyl terminus and, optionally, having amino acid replacements suggested to have improved stability while retaining activity in Eur. Pat. Ap. Pub. No. 326,907 to Seno, et al.; mutants lacking a substantial part of the amino-or carboxyl-terminus such as those described by Seno, et al., *Eur. J. Biochem.* 188: 239-245 (1990); muteins having various point mutations or an N-terminal deletion suggested in Eur. Pat. Ap. Pub. No. 298,723 to Fiddes, et al.; the M1-bFGF to M6-bFGF muteins containing missing and substituted amino acids disclosed in Eur. Pat. Ap. Pub. No. 363,675 to Bergonzoni, cited above; readily expressed FGF prepared by replacement of Ala-3 and Ser-5 of recombinant FGF with Glu as described in Seddon, A.P. et al., *Annals N.Y. Acad. Sci.* 638: 98-108 (1991) and analogues thereof; and the like.

In the practice of this invention, a fibroblast growth factor derivative of this invention such as a human FGF-2 structural analogue is prepared by deleting, adding to, or substituting at least one amino acid, preferably more than one amino acid, in the sequence between residues 112 (Tyr) and 128 (Lys). In some embodiments, the substitution occurs between residues 115 (Tyr) and 124 (Tyr); in others, the change occurs between 118 (Arg) and 122 (Ser). As defined above, the numbering convention for other FGF analogues is relative to FGF-2.

In some preferred embodiments, fibroblast growth factor structural analogues have an amino acid sequence replacement in the ninth or tenth β -strand of the factor, or the sequence that corresponds to the surface loop that connects the ninth and tenth β -strands. Amino acid se-

- 18 -

quences are herein defined as a sequence of at least three amino acids, typically at least five amino acids. Exemplary amino acid sequences are set out above.

For example, a fibroblast growth factor structural analogue such as a human FGF-2 structural analogue is prepared by replacing all or part of the amino acid sequence in the 9th to 10th β -strand surface loop with a random amino acid sequence of any length, provided that the overall folding of the molecule is not significantly perturbed. Thus, a sequence derived from a structurally related polypeptide such as fibroblast growth factor-2 from another species, human fibroblast growth factor-1, or 3 to 9, fibroblast growth factor-1, or 3 to 9 from another species, human interleukin-1 α or β , or interleukin-1 α or β from another species, soybean trypsin inhibitor or hisactophilin, may be employed. In some embodiments, preferred FGF-2 analogues involve an amino acid replacement that eliminates phosphorylation at Thr121.

In one embodiment, a structural analogue of this invention exhibiting desirable biological properties more particularly described below comprises a fibroblast growth factor-2 derivative having surface loop amino acids Arg118-Lys119-Tyr120-Thr121-Ser122 (amino acids 118-122 of SEQ ID NO 1) replaced with corresponding amino acid sequence Ala115-Gln116-Phe117-Pro118-Asn119 (amino acids 231-235 of SEQ ID NO 3) from human interleukin-1 β , denoted FGF-2LI in the Examples that follow (SEQ ID NO 15). In another embodiment, the analogue comprises a human fibroblast growth factor-2 derivative having surface loop amino acids Arg118-Lys119-Tyr120-Thr121-Ser122 (amino acids 118-122 of SEQ ID NO 1) replaced with corresponding amino acid sequence Lys115-Lys116-His117-Ala118-Glu119-Lys120-His121 (amino acids 115-121 of SEQ ID NO 2) derived from bovine FGF-1, denoted FGF-2LA in the Examples that follow (SEQ ID NO 16). In yet another embodi-

- 19 -

ment, the analogue comprises a human fibroblast growth factor-2 derivative having surface loop amino acids 118 to 122 replaced with a corresponding 9-residue amino acid loop sequence from FGF-7 (also known as Keratinocyte Growth Factor), Ala-Lys-Trp-Thr-His-Asn-Gly-Gly-Glu from FGF-7 (residues 154-162 of SEQ ID NO 8).

The novel fibroblast growth factor structural analogues of this invention are prepared by point mutations, sequence alterations or polypeptide assembly from constituent amino acids or peptides using chemical, biochemical or physical means known to those skilled in the art. Alternatively, the novel fibroblast growth factor analogues of this invention are prepared by recombinant protein synthesis involving preparation of DNA encoding a surface loop mutein, insertion of that DNA into a vector, expression of the vector in host cells, and isolation of the mutant FGF thereby produced.

DNA encoding the FGF structural analogues of this invention are prepared by altering a gene of fibroblast growth factor by nucleotide deletions, nucleotide additions, or point mutations produced using standard means. Illustrations are set out in Examples 1 and 2. Because of the degeneracy of the genetic code, a variety of codon change combinations can be selected to form DNA that encodes the FGF analogues of this invention, so that any nucleotide deletion(s), addition(s), or point mutation(s) that result in a DNA encoding loop mutant FGF are encompassed by this invention. Since certain codons are more efficient for polypeptide expression in certain types of organisms, the selection of fibroblast gene alterations to yield DNA material that codes for the FGF muteins of this invention are preferably those that yield the most efficient expression in the type of organism which is to serve as the host of the recombinant vector. Altered

- 20 -

codon selection may also depend upon vector construction considerations.

Fibroblast growth factor DNA starting material which is altered to form DNA coding for the FGF analogues of the invention may be natural, recombinant or synthetic. Thus, DNA starting material is isolated from tissue or tissue culture, constructed from oligonucleotides using conventional methods, obtained commercially, or prepared by isolating RNA coding for FGF from fibroblasts, and using this RNA to synthesize single-stranded cDNA which is used as a template to synthesize the corresponding double stranded DNA.

Illustrating the present invention are cloned complementary DNA sequences defining human FGF-2 analogues such as that constructed in Examples 1 and 2. Also encompassed are DNA sequences homologous or closely related to complementary DNA described herein, namely DNA sequences which hybridize, particularly under stringent conditions that result in pairing only between nucleic acid fragments that have a high frequency of complementary base sequences, to FGF analogue cDNA, and RNA corresponding thereto. In addition to the FGF-encoding sequences, DNA encompassed by this invention may contain additional sequences, depending upon vector construction sequences, that facilitate expression of the gene.

DNA encoding the FGF analogues of this invention, or RNA corresponding thereto, are then inserted into a vector, e.g., a pBR, pUC, pUB or pET series plasmid, and the recombinant vector used to transform a microbial host organisms. Host organisms useful in the invention are bacterial (e.g., *E. coli* or *B. subtilis*), yeast (e.g., *S. cerevisiae*), mammalian (e.g., mouse fibroblast), or insect cells. This invention thus also provides novel, biologically functional viral and circular plasmid RNA and DNA

- 21 -

vectors incorporating RNA and DNA sequences describing the FGF analogues generated by standard means. Culture of host organisms stably transformed or transfected with such vectors under conditions facilitative of large scale
5 expression of the exogenous, vector-borne DNA or RNA sequences and isolation of the desired polypeptides from the growth medium, cellular lysates, or cellular membrane fractions yields the desired products. An example of expression of FGF-2 muteins in *E. coli* is given in Exam-
10 ple 3.

The present invention provides for the total and/or partial manufacture of DNA sequences coding for FGF-2 loop mutants, and including such advantageous characteristics as incorporation of codons preferred for expres-
15 sion by selected non-mammalian hosts, provision of sites of cleavage by restriction by endonuclease enzymes, and provision of additional initial, terminal or intermediate DNA sequences which facilitate construction of readily expressed vectors. Correspondingly, the present inven-
20 tion provides for manufacture (and development by site specific mutagenesis of cDNA and genomic DNA) of DNA sequences coding for microbial expression of FGF analogues which differ from the forms specifically described herein in terms of identity or location of one or more
25 amino acid residues (i.e., deletion analogues containing less than all of the residues specified for human FGF-2, and/or substitution analogues wherein one or more residues are added to a terminal or medial portion of the polypeptide), and which share the biological properties
30 of FGF-2 analogues described herein.

DNA (and RNA) sequences of this invention code for all sequences useful in securing expression in procary-
otic or eucaryotic host cells of polypeptide products having at least a part of the primary structural confor-
35 mation, and on or more of the biological properties of

- 22 -

FGF analogues which are comprehended by: (a) the DNA sequences encoding FGF-2 loop muteins as described herein, or complementary strands; (b) DNA sequences which hybridize to DNA sequences defined in (a) or fragments thereof; and (c) DNA sequences which, but for the degeneracy of the genetic code, would hybridize to the DNA sequences defined in (a) and (b) above. Specifically comprehended are genomic DNA sequences encoding allelic variant forms of FGF analogues included therein, and sequences encoding loop mutein RNA, fragments thereof, and analogues wherein RNA or DNA sequences may incorporate codons facilitating transcription or RNA replication of messenger RNA in non-vertebrate hosts.

Isolation and purification of microbially expressed polypeptides provided by the invention are by conventional means including, for example, preparative chromatographic separations such as that illustrated Example 3, and immunological separations, including monoclonal and/or polyclonal antibody preparations.

As summarized above and described in detail in the Examples below, two example FGF-2 structural analogues of this invention are FGF-2LA (SEQ ID NO 16) and FGF-2LI (SEQ ID NO 15), FGF-2 sequences having loop residues 118 to 122 replaced by corresponding sequences from bovine FGF-1 and human interleukin-1 β , respectively, expressed in *E. coli*. These mutations have no apparent effect on either heparin binding, estimated by the concentration of NaCl required to elute the protein from heparin Sepharose, or on the ability of the proteins to compete with ¹²⁵I-FGF-2 for binding to the FGF receptors present on baby hamster kidney cells and NIH 3T3 cells (Example 4). The loop analogues are mitogenically active in bovine endothelial cell proliferation assays. Both analogues exhibit significantly reduced capability to induce urokinase-typ plasminogen activator. In an *in vitro* angio-

- 23 -

genesis assay, FGF-2LA exhibits capillary-like tube formation comparable to wild-type FGF-2. In the same *in vitro* assay, FGF-2LI exhibits much less induction of capillary-like structures, but it significantly stimulates angiogenesis in an *in vivo* assay.

Another FGF-2 loop mutant, FGF-2LK (described in greater detail in the Examples), contains a 9-amino acid sequence from FGF-7 (Keratinocyte Growth Factor). FGF-2 does not bind to FGF-7 receptor (FGFR2 IIIb) on keratinocytes, and FGF-7 does not bind to FGF receptor type 1 (FGFR1). Replacement of the loop sequence in FGF-2 with that from FGF-7 results in an apparent decrease in the affinity of the protein to heparin. In receptor binding experiments, the FGF-2LK protein is unable to displace or compete with binding of FGF-2 to FGF receptor type 1, but competes with FGF-7 to receptor type 2 (IIIb) whereas FGF-2 does not compete with binding. Thus, the loop sequence confers receptor-ligand specificity and allows for the binding of FGF-2LK mutant to a receptor subtype (FGFR IIIb) that binds FGF-7 but not FGF-2. Conversely, the loop sequence from FGF-7 abolishes binding of the protein to FGF receptor 1.

Introduction of new primary structural elements in the FGF surface loop without significantly perturbing the overall secondary and tertiary structure of the molecule provides an array of FGF structural analogues that give rise to altered properties from one FGF type to another, and thus a means to modulate the activities of FGF proteins, a means to pharmacologically dissociate the biological activities of the proteins, and a means to introduce new activities. As illustrated in the Examples that follow, FGF loop analogues can be structured to bind to the same receptors as corresponding native FGF, or to different receptors, particularly to receptors corresponding to the loop rather than to the native FGF.

- 24 -

FGF antagonists exhibiting reduced biological activity are useful as anticancer and antiproliferative agents. The antagonists that act as angiogenesis inhibitors are useful for the treatment of diseases where neovascularization is dominant in the pathology such as retinopathies of the eye, neovascular glaucoma, skin disorders, chronic inflammation, rheumatoid arthritis, and the like.

FGF loop structural analogues that are agonists of FGF activity can promote vascularization, cell growth, and/or cell survival, and thus have application in tissue repair such as healing of wounds, burns, bone fractures, surgical abrasions, gastrointestinal ulcers, and the like as well as tissue repair during ischemia and myocardial infarction via neovascularization of ischemic tissue.

In addition to surface loop amino acid sequence replacements, this invention further provides growth factor peptide antagonists constructed to mimic the replaced loop, i.e., that bind to FGF receptors but do not stimulate proliferation or migration of fibroblasts and other cells stimulated by the corresponding factors containing the loop.

Expression of FGF receptor type is cell-specific, and the type of receptor expressed determines which FGF the cell will respond to. As mentioned above, change in receptor specificity, such as that seen for FGF-2LA, indicates that FGF-2 can be engineered to target a cell type that it normally does not interact with, such as an epithelial cell rather than an endothelial cell.

The following examples are presented to further illustrate and explain the present invention and should not be taken as limiting in any regard.

- 25 -

Example 1

This example illustrates the construction of a human FGF-2 structural analogue denoted FGF-2LI (SEQ ID NO 15) having the surface loop (residues 118 to 122) of FGF-2 replaced with the corresponding amino acids derived from interleukin-1 β (IL-1 β).

A gene encoding human glu^{3,5}FGF-2 is first prepared as described in Seddon, A.P. et al., cited above, herein incorporated in its entirety by reference, and cloned into a T7 expression vector, pET-3a(M13). Briefly stated, a synthetic gene encoding the 155 amino acid form of human FGF-2 cloned into pUC 18 is purchased from British Bio-technology, Oxford, UK. The nucleotide sequence (2-49) to be changed is excised from pUC 18 with *Hind*III and *Bsp*MII and a synthetic fragment encoding the first 5 N-terminal amino acids of FGF-1 and containing an internal *Nde*I site is cloned into pUC 18, yielding a construct encoding glutamic acid at positions 3 and 5. The cDNA encoding FGF-2 is then excised from pUC 18 with *Nde*I and *Bam*HI and cloned into the *Nde*I and *Bam*HI restriction endonuclease sites of the expression vector pET-3a(M13), a derivative of pET-3a.

Two unique restriction endonuclease sites, *Bst*BI and *Spl*I, are introduced into the gene in such a way as to produce no change in the encoded amino acids (i.e., silent mutations) at positions that flank the codons encoding the segment Ser117-Trp123 of FGF-2 (Figure 1).

Replacement of residues Arg118-Lys119-Tyr120-Thr121-Ser122 of FGF-2 (amino acids 118-122 of SEQ ID NO 1) with the human sequence Ala115-Gln116-Phe117-Pro118-Asn119 (amino acids 231-235 of SEQ ID NO 3) from the corresponding loop of the structural analogue IL-1 β (115-119) (Figure 1) is then effected. The plasmid DNA is subjected to

- 26 -

*Bst*B1 and *Spl*1 digestion and the larger DNA fragment, isolated using agarose gel electrophoresis. The DNA fragment is ligated using T4 DNA ligase to a double-stranded DNA obtained by annealing two synthetic oligonucleotides, 5'-CGAACGATTG GAATCTAATA ACTACAATAC GTACCGGTCT GCGCAGTTTC CTAAGTGGTA TGTGGCACTT AAGC-3' (SEQ ID NO 11) and 5'-GTACGCTTAA GTGCCACATA CCAGTTAGGA AACTGCGCAG ACCGGTACGT ATTGTAGTTA TTAGATTCCA ATCGTT-3' (SEQ ID NO 12), that contain termini compatible to those generated by *Bst*B1 and *Spl*1 digestion. The ligation product is used to transform strain DH5α *E. coli* cells. The desired mutant plasmid is selected for on the basis of susceptibility to cleavage at the newly introduced *Afl*2 restriction site (underlined) and confirmed by complete sequencing of the gene. The plasmid in *E. coli* was deposited in the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852 USA and bears A.T.C.C. accession number 69417.

Example 2

This example illustrates the construction of a human FGF-2 structural analogue denoted FGF-2LA (SEQ ID NO 16) having the surface loop (residues 118 to 122) of FGF-2 replaced with the corresponding amino acid sequence derived from FGF-1 (Figure 1).

Replacement of the segment Arg118-Ser122 of FGF-2 with the bovine sequence Lys115-Lys116-His117-Ala118-Glu119-Lys120-His121 (amino acids 115-121 of SEQ ID NO 2; Asn in the human sequence) corresponding to the surface loop 115-121 of FGF-1 (see Figure 1) is accomplished as described in Example 1 above using the following annealed synthetic mutagenic oligonucleotides: 5'-CGAACGATTG GAATCTAATA ACTACAATAC GTACCGGTCT AAAAAGCATG CTGAAAAACA CTGGTATGTG GCACTTAAGC-3' (SEQ ID NO 13) AND 5'-GTACGCTTAA GTGCCACATA CCAGTGT TTTT TCAGCATGCT TTTTAGACCG GTACGTATTG

- 27 -

TAGTTATTAG ATTCCAATCG TT-3' (SEQ ID NO 14). The desired mutant plasmid is selected on the basis of the susceptibility to cleavage at the newly introduced *Sph*I restriction site (underlined) and confirmed by complete sequencing of the gene.

Example 3

FGF-2LA and FGF-2LI mutants constructed in Examples 1 and 2 are expressed and purified in this Example.

Following sequence verification, the plasmids containing FGF-2 loop mutants described in Examples 1 and 2 above are transformed into competent *E. coli* BL21 plys S and cultured at 37°C in Luria broth containing 50 µg/ml ampicillin and 30 µg/ml chloramphenicol until an absorbance at 600 nm of 0.4 is reached. Expression of the recombinant protein is induced by the addition of 2 mM isopropylthiogalactoside for 2 hours at 37°C.

Cells from 1 liter cultures are harvested by centrifugation, resuspended in 30 ml of 50 mM Tris-HCl, pH 7.5 containing 0.1 mM EDTA and 0.6 M NaCl, and disrupted by treatment with lysozyme (10 µg/ml) for 20 min at 4°C followed by sonication (6 x 30 sec pulses). The lysates are clarified by centrifugation (10,000 x g; 20 min) and the supernatant solutions incubated with 5 ml of hydrated heparin Sepharose (Pharmacia/LKB) at 4°C for 1 hour with constant rotation. The resin is isolated by filtration on 0.8 µm filter apparatus (Nalgene), washed extensively with 10 mM Tris-HCl pH 7.4 containing 0.6 M NaCl and bound protein eluted with Tris buffer containing 3 M NaCl (25 ml). The 3 M NaCl eluent is diluted 6-fold with Tris buffer and loaded onto a TSK heparin 5PW column (0.21 x 15 cm; The Nest Group, MA) and the column developed using a linear NaCl gradient (0.6 to 2 M) in 90 min at a flow rate of 3 ml/min.

Not furnished upon filing

- 29 -

residues. The proteins exhibit 36 other sites of identical and conserved residues in addition to the 7 denoted.

Example 4

5 This example describes binding and cell proliferation studies using the FGF-2 mutants isolated and purified in Example 3 above.

The affinities of FGF-2LA and FGF-2LI for immobilized heparin are identical to wild type FGF-2 on elution from a TSK-heparin column at about 1.5 M NaCl.

10 FGF-2LI and FGF-2LA are tested for their capacity to compete for the binding of ^{125}I -FGF-2 (Amersham Corp.) to baby hamster kidney (BHK) cells, which express high numbers of FGF receptor. The assay employed is described by Moscatelli (*J. Cell. Physiol.* 131: 123-130 (1987)).
15 Briefly stated, BHK cells plated on 24-well plates are incubated with 50 pM ^{125}I -FGF-2 with serial dilutions of unlabelled FGF-2, FGF-2LI or FGF-2LA loop mutants at room temperature for 1 hour or at 4°C for 2 hours. The cells are then incubated at 4°C for 30 minutes, washed twice
20 with phosphate-buffered saline and treated with 20 mM HEPES (N-2-hydroxyethyl-piperazine-N'-2-ethanesulfonic acid), pH 7.5, containing 2 M NaCl to remove ^{125}I -FGF-2 bound to low affinity heparin sulfate binding sites. Receptor bound ^{125}I -FGF-2 is recovered by treatment of the
25 cells with 0.5% Triton X-100 in 0.1 M sodium phosphate, pH 8, and counted in a γ -counter. Assays are conducted in duplicate.

The results of this binding assay, plotted in Figure 2, are found to be similar for FGF-2, FGF-2LA and FGF-
30 2LI. Replacement of FGF-2 sequence 118-122 with the corresponding IL-1 β or FGF-1 sequences has no apparent effect on the capacities of the mutant proteins to com-

- 30 -

pete with ^{125}I -FGF-2 binding to high affinity cell membrane FGF type I receptors present on BHK cells. Since the mutants contain FGF-1 and IL-1 β sequences, IL-1 β and FGF-1 are tested in the BHK cell FGF-receptor binding assay. FGF-1 binds to BHK cell FGF receptors with an affinity equal to that of FGF-2, whereas no binding for hrIL-1 β (Biogen) is detected.

Figure 3 shows the binding of radioidinated FGF-2 (bFGF), FGF-LA and FGF-LI to FGF-receptor type 1 present on NIH 3T3 cells as a function of added unlabelled FGF proteins using these procedures. The data show that the competition binding curves for FGF-LA and FGF-LI are identical to that for FGF-2, and demonstrate that the loop exchanges in FGF-2 have no impact on the receptor binding properties of the proteins to cell surface FGF receptor type 1 (FGFR1).

The mitogenic activity of FGF-2, FGF-2LI and FGF-2LA are determined using bovine vascular endothelial cells derived from adult aortic arch as previously described (Gospardarowicz, D., et al., *Proc. Nat. Acad. Sci.* 81: 6963-6967 (1984)). Briefly, cells are seeded at an initial density of 0.8×10^4 cells per 24-well plate in 0.5 ml Dulbecco's modified Eagle's medium (DMEM) containing 10% calf serum (Hyclone, Logan, UT) supplemented with penicillin (100 units/ml), streptomycin (100 ug/ml) and L-glutamine (2 mM). Two hours after plating, 20 ul aliquots of serial dilutions of FGF-2 and the mutants in DMEM are added. Inhibition of FGF-2-stimulated growth was determined at a fixed concentration of FGF-2, while that of the mutant proteins is varied. After 5 days in culture, duplicate plates are trypsinized and cell densities determined by cell counting in a Coulter counter. Determinations are conducted in duplicate.

- 31 -

The results of this cell proliferation assay are plotted in Figure 4. The dashed line indicates basal cell growth in the absence of added FGF-2. Closed circles indicate stimulation by FGF-2. FGF-2LI mutant is represented by open squares and FGF-2LA mutant by open triangles. FGF-2LI is about 5 to 10 times less potent than FGF-2, whereas FGF-2LA is as potent as the wild-type factor.

Example 5

10 This example describes *in vitro* and *in vivo* angiogenesis studies using the FGF-2 mutants isolated and purified in Example 3 above.

Urokinase-type plasminogen activator induction by the FGF mutants is evaluated. Adult bovine aortic endothelial (ABAE) cells are seeded at 20,000 cells/well in 96-well plates and maintained in DMEM containing 10% calf serum (Hyclone, Logan, UT) supplemented with penicillin (100 units/ml), streptomycin (100 µg/ml) and L-glutamine (2 mM) and different concentrations of FGF-2 or mutants. After 24 hours, the cells are washed with phosphate-buffered saline and lysed in 60 mM Tris-HCl, pH 8.5, containing 0.05% Triton® X-100. Cell-associated urokinase-type plasminogen activator (uPA) activity is measured as described by Presta, et al., cited above, using the plasmin chromogenic substrate D-norleucyl-hexahydrotyrosyl-lysine p-nitroanilide acetate (American Diagnostics, Greenwich, CT). Cell-associated protein concentrations are determined using Coomassie blue binding to protein. The results are plotted in Figure 5. Both FGF-2LI and FGF-2LA exhibit significantly reduced capabilities to induce urokinase-type plasminogen activator.

In vitro angiogenesis evaluations are made by observing whether the mutants induce capillary-like struc-

- 32 -

tures in ABAE cells cultured on a 3-dimensional collagen gel. Three-dimensional collagen gel plates (24-well) are prepared by adding 0.5 ml chilled solution of 0.7 mg/ml rat-tail type I collagen (Becton Dickinson Labwares, Bedford, MA) containing DMEM and adjusting to neutral pH with NaHCO_3 to each well. After formation of collagen gel (about 1-2 mm thickness), ABAE cells are seeded at 50,000 cells/well. The cultures are maintained at 37°C in DMEM containing 10% calf serum (Hyclone, Logan, UT) supplemented with penicillin (100 units/ml), streptomycin (100 $\mu\text{g}/\text{ml}$) and L-glutamine (2 mM) until the cultures reach confluency, usually in 5 days by which time the cells form a monolayer on the gel. The medium is then replaced with fresh medium containing different concentrations of FGF-2 or mutants. The cultures are maintained at 37°C for 48 hours, then discontinued by fixation with cold methanol (-20°C).

The abundance of the capillary-like structures formed by ABAE cells is analyzed by using a Kontron IBAS Image Analyzer assisted with a Hamamatsu C2400 video camera and a Zeiss Axioshop microscope. The phase-contrast image of each field obtained with a video camera is converted to a binary image in which the areas occupied by a capillary-like structure is white and the rest is a black background. The extent of *in vitro* angiogenesis is then measured as a percentage of the white areas. Cell cultures of identical conditions are maintained in duplicate wells. Three to four areas of each well are examined by image analysis and the mean value and standard deviation are determined. The results from this computer-assisted quantitation method are set out in Figure 6. The induction of tube formation by mutant FGF-2LI is much less than that by wild-type FGF-2, whereas that by FGF-2LA is comparable with the wildtype.

- 33 -

In an *in vivo* model of angiogenesis, FGF-2LI is evaluated using the rabbit ear chamber system modified and improved by Howden, G.F., and Silver, I.A., (*Int. Endodontic J.* 13: 3-16 (1980)), observing the growth
5 factor-induced neovascularization, or its inhibition, by modified growth factors or other reagents.

Rabbits are sedated using 50-70 mg/kg ketamine and 10 mg/kg xylozine administered intramuscularly. The ears are shaved with an electric hair clipper and cleaned with
10 water, followed by Betadine®. To minimize the risk of infection, the rabbits are given 20,000 units/kg benzathine penicillin intramuscularly prior to insertion of the ear chambers. The chamber is inserted into an approximately 0.75 cm area which is not crossed by any
15 major vessels. During the course of the experiments, observations or any minor handling of ear chambers of the animals is done by sedating the animal using the intramuscular administration of a combination of Butorphanol® and acepromazine at levels of 1 mg/kg each. FGF-2 or
20 FGF-2LI bound to heparin-Sepharose® beads in a controlled-release alginate capsule is contained in one of the ear chambers and the other is used as a control containing only the heparin-Sepharose® alginate capsule vehicle. Thus, the effect of the growth factors and the
25 controlled-release capsule alone can be observed simultaneously in the same animal. FGF-2 and FGF-2LI are tested at 1, 10, and 100 ng/chamber in triplicate. The animals are kept for 4 to 6 weeks and neovascularization is observed visually and recorded photographically at weekly
30 intervals.

The criteria for evaluation of the degree of neovascularization using this procedure is set out below.

- 34 -

Criteria for Evaluation of the Degree of Neovascularization

	Observation	Score
	No Vascularization	0
	Small Buds	0.5
5	Extensive Buds	1.0
	Capillary Network Extending to $\frac{1}{2}$ of the Chamber	2.0
	Capillary Network Extending $> \frac{1}{2}$ of the Chamber; No Anastomoses	3.0
	Anastomoses from Opposite Sides of the Chamber	4.0
	Extensive Anastomosing Capillary Network Filling Chamber Area	5.0

- 10 The data, plotted in Figure 9, show that FGF-2LI is superior to FGF-2 in its potential to induce angiogenesis. Both FGF-2 and FGF-2LI stimulate large vessel formation, their growth across the chamber, resulting in anastomosing vessels with an active blood flow. In comparison,
- 15 in control chambers that contain alginate-heparin Sepharose® beads, void of exogeneously added growth factors, only short capillaries are formed. The difference in the activity of FGF-2LI compared to FGF-2 is both quantitative and qualitative. Striking stimulatory activity of FGF-2LI compared to FGF-2 is observed over a
- 20 wide range of concentrations (i.e., from 0.1 - 400 ng/chamber), and the stimulation is prolific and sustained beyond the time when the stimulatory effect of FGF-2 has subsided. (See Figure 8.)
- 25 Though FGF-2 shows a dose-dependent induction of new blood vessels compared to the vehicle alone, a comparable activity for FGF-2 is observed only at concentrations of 10-100 ng/chamber; at higher dosages, a diminished response is observed. Interestingly, at 400 ng/chamber of
- 30 FGF-2, angiogenesis is completely inhibited, while FGF-2LI at the same dose greatly stimulates formation of capillaries and larger vessels. Thus, there is an observed difference in the biological activity of FGF-2LI compared to FGF-2.
- 35 Although the *in vitro* data suggest a reduced angiogenic activity for FGF-2LI, the *in vivo* studies indicate

WO 95/08630

-35-

PCT/US94/10800

Not furnished upon filing

- 36 -

Labware Division, Becton Dickinson (Oxnard, CA). Four-well tissue culture plates are from Nunc (Roskilde, Denmark).

Soluble FGF receptor proteins are constructed by cloning of the extracellular region of murine FGF receptor 1 (FGFR-1; flg), FGF receptor 2 (FGFR-2; bek) or the KGF receptor (FGFR(IIIb); K-sam) into the alkaline phosphatase-tag expression vector, which encodes for a secreted form of placental alkaline phosphatase (AP). The FGF receptor alkaline phosphatase (FRAP) plasmids are cotransfected into NIH 3T3 cells by electroporation with a selectable neomycin resistance gene. Clonies are selected in G418 (600 $\mu\text{g}/\text{ml}$) and screened for secreted AP enzyme activity in the conditioned medium. Clones of each receptor which produced a high level of AP activity (2 to 4 A_{405} units/min/ml) are then used to produce conditioned medium for binding assays.

Components of the soluble receptor binding reaction mixture include FRAP-conditioned medium (0.24 OD units/min), 2 ng/ml ^{125}I -FGFs and 200 ng/ml heparin. The FGF:heparin:FRAP ternary complex is immunoprecipitated with 20 μl of a 1:1 slurry of anti-AP monoclonal antibodies coupled to protein A Sepharose®. All components are mixed at room temperature. The total volume is adjusted to 200 μl by addition of DMEM containing 0.1% bovine serum albumin. Binding is allowed to proceed for 1 to 2 hours at 24°C, after which time bound receptor complex or the ligand is recovered by centrifugation at 4°C (10 s at 2,000 x g). The pelleted material is washed twice with 500 μl of an ice cold buffer containing HEPES (20 mM), NaCl (150 mM), glycerol (10%) and Triton® X-100 (1%). ^{125}I -FGF binding is quantitated by counting of the samples in a gamma counter (LKB). Alternatively, AP enzyme activity of the FRAP protein is determined by transferring the FRAP receptor bound to heparin-Sepharose® to a flat-

- 37 -

bottom microtiter plate in a volume of 50 μ l of PBS. The reaction is initiated by addition of substrate (50 μ l of 2x solution of AP assay buffer containing 2 M diethanolamine, 1 mM $MgCl_2$, 20 mM homoarginine and 12 mM p-nitrophenyl phosphate). The reaction is followed at room temperature at 405 nm in a kinetic microplate reader.

Receptor binding is determined by quantitating release of labelled FGF from receptors. Briefly, FGF bound to heparan sulfate low affinity sites is released from the cell surface by a 5 minute incubation with an ice cold solution containing 1.6 M NaCl, 20 mM HEPES, pH 7.4, and the amount of radioactivity release determined in a gamma-counter. FGF bound to high affinity receptors is dissociated by a 2 M NaCl (20 mM acetate buffer, pH 4.0) extraction, and the released labelled FGF is quantitated.

Chemical cross-linking experiments are carried out at room temperature in a volume of 20 μ l in siliconized 0.5-ml microcentrifuge tubes. The reaction mixtures contain FGF receptor immobilized to anti-AP monoclonal antibodies coupled to protein A Sepharose®, 200 ng/ml heparin, 2 ng/ml ^{125}I -bFGF, 20 mM phosphate buffer (pH 7.4), and 140 mM NaCl. After a 90 minute incubation, 1 ml of a solution of disuccinimidyl suberate (Pierce) dissolved in dimethyl sulfoxide is added to give a final concentration of 0.15 mM, and the mixture incubated for an additional 30 minutes. The reaction is quenched by addition of 1 ml of 200 mM ethanolamine-HCl (pH 8.0) for 30 min. The reaction mixtures are diluted 1:1 with 2x SDS-polyacrylamide gel electrophoresis loading buffer and electrophoresed on an SDS-12% polyacrylamide gel. Cross-linked FGF to the FGF receptor are detected by autoradiography on Kodak XAR film.

The position of the surface loop in FGF-2 coincides with a variable sequence region in the FGF family of

- 38 -

proteins where various insertions occur. Since this region may be involved with determining ligand-receptor binding specificity the binding profiles of the loop mutants, LA and LI, to FGFR1 (Flg) and FGFR2 sub-type IIIb (FGF-7 or KGF receptor), which does not bind FGF-2 are determined. Figures 9A and 9B show the competition binding curves to soluble versions of FGFR1 and FGFR2(IIIb) for FGF-1, FGF-2, FGF-7 and the loop mutants following the displacement of radioiodinated FGF-2 and FGF-7 from the soluble receptor, respectively. FGF-1, FGF-2, FGF-2LA and FGF-2LI bind equally well to FGFR1, but no binding of FGF-7 is detected (Figure 9A).

The binding profiles for the binding of various FGF proteins to FGFR2(IIIb) (Figure 9B) show that FGF-7 and FGF-1 bind with the same affinity and that FGF-2 and the FGF-2LI mutant do not bind to this receptor type; however, the binding of the LA mutant to FGFR2(IIIb) is only about 5-times less than that of FGF-7 or FGF-1. That the binding profile of FGF-2LA now mirrors that for FGF-1 and not of FGF-2, the FGF-1 loop-host protein, is consistent with the involvement of this surface loop in determining receptor-ligand specificity.

The binding specificities of radiolabelled FGF proteins and loop mutants to various soluble FGF receptors are confirmed by chemical cross-linking of the factors to the receptors in the presence and absence of an excess of the unlabelled FGF. FGF-1, FGF-2, FGF-2LA and FGF-2LI bind to FGFR1, and addition of an excess of the unlabelled FGF abolished cross-linking of ¹²⁵I-FGF. Cross-linking of ¹²⁵I-FGF-7 to FGFR1 is not detected. An identical cross-linking profile is obtained when the binding experiments are repeated using a soluble FGFR2 receptor. The cross-linking profile for FGFR2(IIIb) reveal that only FGF-1, FGF-7 and FGF-2LA bind to this receptor and that FGF-2 and the FGF-2LI mutant are excluded from bind-

- 39 -

ing. The cross-linking studies are consistent with the binding data presented in Figures 2, 3, and 8.

Example 7

5 The FGF-7 loop sequence is introduced into FGF-2, and the properties of the new FGF-2 loop mutant are observed in this example.

A FGF-2 loop mutant denoted FGF-2LK containing the corresponding 9-residue loop sequence Ala-Lys-Trp-Thr-His-Asn-Gly-Gly-Glu from FGF-7 (residues 154-162 of SEQ ID NO 8) is constructed using the procedures outlined in Examples 1 and 2 above.

15 FGF-2LK shows an apparent decrease in affinity for heparin. NaCl elution from heparin-Sepharose® is about 0.7 M NaCl, compared with 1.4 M NaCl for the wild-type protein. The molarity of NaCl required to elute the LK mutant is similar to that required to elute FGF-7. The lowered affinity of the FGF-2LK mutant for heparin suggests that the loop sequence, although not directly involved in binding to heparin, is able to modify the affinity of the protein for heparin.

25 In receptor binding experiments like those set out in Example 6 above, the FGF-2LK protein is unable to displace or compete with the binding of ¹²⁵I-FGF-2 to FGF receptor type 1, but is able to compete with binding of ¹²⁵I-FGF-7 to FGF receptor type 2 (IIIb), whereas FGF-2 does not compete with binding. Although the potency of the FGF-2LK is about 100 times weaker than the competition observed using unlabelled FGF-7, a clear change in receptor-ligand binding profiles is observed.

30 The results indicate that the loop sequence from FGF-7 confers receptor-ligand specificity, and allows for

- 40 -

the binding of the FGF-2LK mutant to a receptor subtype that binds FGF-7 but not FGF-2. Conversely, the loop sequence from FGF-7 in the FGF-2 host molecule abolishes binding of the protein to FGF receptor 1. That the affinity of the interaction is decreased by a factor of 100 suggests that other determinants in FGF-7 are involved in contributing to the binding affinity of the ligand to the receptor. The evidence also suggests that the loop sequences may modify binding to heparin.

10 The above description is for the purpose of teaching the person of ordinary skill in the art how to practice the present invention, and it is not intended to detail all those obvious modifications and variations of it which will become apparent to the skilled worker upon
15 reading the description. It is intended, however, that all such obvious modifications and variations be included within the scope of the present invention as defined in the appended claims.

- 41 -

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANTS: Andrew P. Seddon
Luyuan Li
Peter Böhlen
Magdalena Eisinger
- (ii) TITLE OF INVENTION: Surface Loop Structural Analogues of Fibroblast Growth Factors
- (iii) NUMBER OF SEQUENCES: 16
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: American Cyanamid Company
Patent Law Department
 - (B) STREET: One Cyanamid Plaza
 - (C) CITY: Fort Wayne
 - (D) STATE: NJ
 - (E) COUNTRY: United States
 - (F) ZIP: 07470-8426
- (v) COMPUTER READABLE FORM
 - (A) MEDIUM TYPE: 3.5" 1.44 mb diskette
 - (B) COMPUTER: IBM PC
 - (C) OPERATING SYSTEM: MS DOS
 - (D) SOFTWARE: Word Processor
- (vi) PRIOR APPLICATION DATA (CIP of)
 - (A) APPLICATION NUMBER: 08/126,973
 - (B) FILING DATE: 09-24-93
- (viii) ATTORNEY INFORMATION
 - (A) NAME: Estelle J. Tsevdos
 - (B) REGISTRATION NUMBER: 31145
 - (C) REFERENCE/DOCKET NUMBER: 854-008CIP (32,063)
- (ix) TELECOMMUNICATION INFORMATION
 - (A) TELEPHONE NUMBER: 201-831-3242
 - (B) TELEFAX NUMBER: 201-831-3305

- 42 -

(2) INFORMATION FOR SEQ ID NO: 1

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 21
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

- (A) NAME: fibroblast growth factor-2 loop region

(x) PUBLICATION INFORMATION

- (A) AUTHOR: Eriksson, A.E., et al.
- (B) TITLE: Three-dimensional structure of human basic fibroblast growth factor
- (C) JOURNAL: *Proc. Nat. Acad. Sci. USA*
- (D) VOLUME: 88
- (F) PAGES: 3441-3445; sequence on page 3444
- (G) DATE: April 1991
- (K) RELEVANT RESIDUES: segment corresponding to polypeptide residues 110 to 130

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 1:

Asn	Asn	Tyr	Asn	Thr	Tyr	Arg	Ser	Arg	Lys	Tyr	Thr	Ser	Trp	Tyr
110					115					120				
Val	Ala	Leu	Lys	Arg	Thr									
125					130									

SUBSTITUTE SHEET (RULE 26)

- 43 -

(3) INFORMATION FOR SEQ ID NO: 2

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 23
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

- (A) NAME: fibroblast growth factor-1 loop
region

(x) PUBLICATION INFORMATION

- (A) AUTHOR: Eriksson, A.E., et al.
- (B) TITLE: Three-dimensional structure of
human basic fibroblast growth factor
- (C) JOURNAL: *Proc. Nat. Acad. Sci. USA*
- (D) VOLUME: 88
- (F) PAGES: 3441-3445; sequence on page 3444
- (G) DATE: April 1991
- (K) RELEVANT RESIDUES: segment corresponding
to polypeptide residues 107 to 129

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 2:

Asn	His	Tyr	Asn	Thr	Tyr	Ile	Ser	Lys	Lys	His	Ala	Glu	Lys	His
			110					115					120	
Trp	Phe	Val	Gly	Leu	Lys	Lys	Asn							
			125											

- 44 -

(4) INFORMATION FOR SEQ ID NO: 3

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 21
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

- (A) NAME: interleukin-1 β loop region

(x) PUBLICATION INFORMATION

- (A) AUTHOR: Eriksson, A.E., et al.
- (B) TITLE: Three-dimensional structure of human basic fibroblast growth factor
- (C) JOURNAL: *Proc. Nat. Acad. Sci. USA*
- (D) VOLUME: 88
- (F) PAGES: 3441-3445; sequence on page 3444
- (G) DATE: April 1991
- (K) RELEVANT RESIDUES: segment corresponding to polypeptide residues 223 to 243

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 3:

Asn	Asn	Lys	Leu	Glu	Phe	Glu	Ser	Ala	Gln	Phe	Pro	Asn	Trp	Tyr
	225						230					235		
Ile	Ser	Thr	Ser	Glu	Ala									
		240												

SUBSTITUTE SHEET (RULE 26)

- 45 -

(5) INFORMATION FOR SEQ ID NO: 4

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 37

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE

(A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

(A) NAME: fibroblast growth factor-3 loop
region

(x) PUBLICATION INFORMATION

(A) AUTHOR: Miyamoto, M., et al.

(B) TITLE (excerpt): Molecular cloning of a
Novel Cytokine cDNA Encoding the Ninth Member
of the Fibroblast Growth Factor Family(C) JOURNAL: *Molecular and Cellular Biology*

(D) VOLUME: 13

(E) NUMBER: 7

(F) PAGES: 4251-4259; Figure 2 on page 4254

(G) DATE: July 1993

(K) RELEVANT RESIDUES: segment corresponding
to polypeptide residues 124 to 160

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 4:

Leu	Gly	Tyr	Asn	Thr	Tyr	Ala	Ser	Arg	Leu	Tyr	Arg	Thr	Val	Ser
125						130					135			
Ser	Thr	Pro	Gly	Ala	Arg	Arg	Gln	Pro	Ser	Ala	Glu	Arg	Leu	Trp
140						145					150			
Tyr	Val	Ser	Val	Asn	Gly	Lys								
155						160								

- 46 -

(6) INFORMATION FOR SEQ ID NO: 5

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 21
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

- (A) NAME: fibroblast growth factor-4 loop
region

(x) PUBLICATION INFORMATION

- (A) AUTHOR: Burgess, W.H., and Maciag, T.
- (B) TITLE: The Heparin-Binding (Fibroblast)
Growth Factor Family of Proteins
- (C) JOURNAL: *Ann. Rev. Biochem.*
- (D) VOLUME: 58
- (F) PAGES: 575-606, Figure 1 on page 580
- (G) DATE: 1989
- (K) RELEVANT RESIDUES: segment corresponding
to polypeptide residues 164 to 184

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 5:

Asn Asn Tyr Asn Ala Tyr Glu Ser Tyr Lys Tyr Pro Gly Met Phe
165 170 175
Ile Ala Leu Ser Lys Asn
180

SUBSTITUTE SHEET (RULE 26)

- 47 -

(7) INFORMATION FOR SEQ ID NO: 6

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 27
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

- (A) NAME: fibroblast growth factor-5 loop
region

(x) PUBLICATION INFORMATION

- (A) AUTHOR: Burgess, W.H., and Maciag, T.
- (B) TITLE: The Heparin-Binding (Fibroblast)
Growth Factor Family of Proteins
- (C) JOURNAL: *Ann. Rev. Biochem.*
- (D) VOLUME: 58
- (F) PAGES: 575-606, Figure 1 on page 580
- (G) DATE: 1989
- (K) RELEVANT RESIDUES: segment corresponding
to polypeptide residues 168 to 194

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 6:

Asn	Ser	Tyr	Asn	Thr	Tyr	Ala	Ser	Ala	Ile	His	Arg	Thr	Glu	Lys
		170					175					180		
Thr	Gly	Arg	Glu	Trp	Tyr	Val	Ala	Leu	Asn	Lys	Arg			
		185					190							

- 48 -

(8) INFORMATION FOR SEQ ID NO: 7

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 21 residues
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

- (A) NAME: fibroblast growth factor-6 loop region

(x) PUBLICATION INFORMATION

(A) AUTHOR: Miyamoto, M., et al.

(B) TITLE (excerpt): Molecular cloning of a Novel Cytokine cDNA Encoding the Ninth Member of the Fibroblast Growth Factor Family

(C) JOURNAL: *Molecular and Cellular Biology*

(D) VOLUME: 13

(E) NUMBER: 7

(F) PAGES: 4251-4259; Figure 2 on page 4254

(G) DATE: July 1993

(K) RELEVANT RESIDUES: segment corresponding to polypeptide residues 166 to 186

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 7:

Asn	Asn	Tyr	Asn	Ala	Tyr	Glu	Ser	Asp	Leu	Tyr	Gln	Gly	Thr	Tyr
				170					175					180
Ile	Ala	Leu	Ser	Lys	Tyr									
				185										

- 49 -

(9) INFORMATION FOR SEQ ID NO: 8

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 25 residues
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

- (A) NAME: fibroblast growth factor-7 loop region

(x) PUBLICATION INFORMATION

- (A) AUTHOR: Miyamoto, M., et al.
- (B) TITLE (excerpt): Molecular cloning of a Novel Cytokine cDNA Encoding the Ninth Member of the Fibroblast Growth Factor Family
- (C) JOURNAL: *Molecular and Cellular Biology*
- (D) VOLUME: 13
- (E) NUMBER: 7
- (F) PAGES: 4251-4259; Figure 2 on page 4254
- (G) DATE: July 1993
- (K) RELEVANT RESIDUES: segment corresponding to polypeptide residues 146 to 170

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 8:

Asn His Tyr Asn Thr Tyr Ala Ser Ala Lys Trp Thr His Asn Gly
 150 155 160

Gly Glu Met Phe Val Ala Leu Asn Gln Lys
 165

- 50 -

(10) INFORMATION FOR SEQ ID NO: 9

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 21 residues
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

- (A) NAME: fibroblast growth factor-8 loop
region

(x) PUBLICATION INFORMATION

- (A) AUTHOR: Tanaka, A., et al.
- (B) TITLE (excerpt): Cloning and Characterization of an Antrogen-Induced Growth Factor
- (C) JOURNAL: *Proc. Natl. Acad. Sci. USA*
- (D) VOLUME: 89
- (F) PAGES: 8928-8931; Figure 2 on page 8930
- (G) DATE: October 1992
- (K) RELEVANT RESIDUES: segment corresponding
to polypeptide residues 136 to 156

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 9:

Asn	Asn	Tyr	Thr	Ala	Leu	Gln	Asn	Ala	Lys	Tyr	Glu	Gly	Trp	Tyr
				140					145					150
Met	Ala	Phe	Thr	Arg	Lys									
				155										

SUBSTITUTE SHEET (RULE 26)

- 51 -

(11) INFORMATION FOR SEQ ID NO: 10

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 27 residues
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: peptide

(v) FRAGMENT TYPE: internal fragment

(ix) FEATURE

- (A) NAME: fibroblast growth factor-9 loop region

(x) PUBLICATION INFORMATION

- (A) AUTHOR: Miyamoto, M., et al.
- (B) TITLE (excerpt): Molecular cloning of a Novel Cytokine cDNA Encoding the Ninth Member of the Fibroblast Growth Factor Family
- (C) JOURNAL: *Molecular and Cellular Biology*
- (D) VOLUME: 13
- (E) NUMBER: 7
- (F) PAGES: 4251-4259; Figure 2 on page 4254
- (G) DATE: July 1993
- (K) RELEVANT RESIDUES: segment corresponding to polypeptide residues 143 to 167

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 10:

Asn	Trp	Tyr	Asn	Thr	Tyr	Ser	Ser	Asn	Leu	Tyr	Lys	His	Val	Asp
		145					150					155		
Thr	Gly	Arg	Arg	Tyr	Tyr	Val	Ala	Leu	Asn	Lys	Asp			
		160					165							

- 52 -

(12) INFORMATION FOR SEQ ID NO: 11

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 74
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: oligonucleotide

(v) FRAGMENT TYPE: synthetic DNA

(ix) FEATURE

- (D) OTHER INFORMATION: used in
preparing constructs

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 11:

CGAACGATTG GAATCTAATA ACTACAATAC GTACCGGTCT GCGCAGTTTC 50
CTAACTGGTA TGTGGCACTT AAGC 74

(13) INFORMATION FOR SEQ ID NO: 12

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 76
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: oligonucleotide

(v) FRAGMENT TYPE: synthetic DNA

(ix) FEATURE

- (D) OTHER INFORMATION: used in
preparing constructs

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 12:

GTACGCTTAA GTGCCACATA CCAGTTAGGA AACTGCGCAG ACCGGTACGT 50
ATTGTAGTTA TTAGATTCCA ATCGTT 76

- 53 -

(14) INFORMATION FOR SEQ ID NO: 13

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 80
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: oligonucleotide

(v) FRAGMENT TYPE: synthetic DNA

(ix) FEATURE

- (D) OTHER INFORMATION: used in
preparing constructs

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 13:

CGAACGATTG GAATCTAATA ACTACAATAC GTACCGGTCT AAAAAGCATG 50
CTGAAAAACA CTGGTATGTG GCACTTAAGC 80

(15) INFORMATION FOR SEQ ID NO: 14

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 82
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: oligonucleotide

(v) FRAGMENT TYPE: synthetic DNA

(ix) FEATURE

- (D) OTHER INFORMATION: used in
preparing constructs

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 14:

GTACGCTTAA GTGCCACATA CCAGTGTTTT TCAGCATGCT TTTTAGACCG 50
GTACGTATTG TAGTTATTAG ATTCCAATCG TT 82

- 54 -

(16) INFORMATION FOR SEQ ID NO: 15

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 471 bases encoding 155 amino acids

(B) TYPE: nucleic acid and amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE

(A) DESCRIPTION: DNA encoding a protein

(v) FRAGMENT TYPE: entire sequence

(vi) IMMEDIATE SOURCE: constructed

(ix) FEATURE

(D) INFORMATION: FGF-2 having surface loop residues 118 to 122 replaced with corresponding structural elements from interleukin-1 β

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 15:

ATG GCT GAA GGG GAA ATC ACC ACG CTG CCC GCC CTT CCG GAG GAT	45
Met Ala Glu Gly Glu Ile Thr Thr Leu Pro Ala Leu Pro Glu Asp	
5 10 15	
GGC GGC AGC GGC GCC TTC CCG CCC GGG CAC TTC AAG GAC CCC AAG	90
Gly Gly Ser Gly Ala Phe Pro Pro Gly His Phe Lys Asp Pro Lys	
20 25 30	
CGG CTG TAC TGC AAA AAC GGG GGC TTC TTC CTG CGC ATC CAC CCC	135
Arg Leu Tyr Cys Lys Asn Gly Gly Phe Phe Leu Arg Ile His Pro	
35 40 45	
GAC GGC CGA GTT GAC GGG GTC CGG GAG AAG AGC GAC CCT CAC ATC	180
Asp Gly Arg Val Asp Gly Val Arg Glu Lys Ser Asp Pro His Ile	
50 55 60	
AAG CTA CAA CTT CAA GCA GAA GAG AGA GGA GTT GTG TCT ATC AAA	225
Lys Leu Gln Leu Gln Ala Glu Glu Arg Gly Val Val Ser Ile Lys	
65 70 75	
GGA GTG TGT GCT AAC CGG TAC CTG GCT ATG AAG GAA GAT GGA AGA	270
Gly Val Cys Ala Asn Arg Tyr Leu Ala Met Lys Glu Asp Gly Arg	
80 85 90	
TTA CTG GCT TCT AAA TGT GTT ACG GAT GAG TGT TTC TTT TTT GAA	315
Leu Leu Ala Ser Lys Cys Val Thr Asp Glu Cys Phe Phe Phe Glu	
95 100 105	

- 55 -

CGA TTG GAA TCT AAT AAC TAC AAT ACT TAC CGG TCT GCG CAG TTT 360
 Arg Leu Glu Ser Asn Asn Tyr Asn Thr Tyr Arg Ser Ala Gln Phe
 110 115 120

CCT AAC TGG TAT GTG GCA TTG AAA CGA ACT GGG CAG TAT AAA CTT 405
 Pro Asn Trp Tyr Val Ala Leu Lys Arg Thr Gly Gln Tyr Lys Leu
 125 130 135

GGT TCC AAA ACA GGA CCT GGG CAG AAA GCT ATA CTT TTT CTT CCA 450
 Gly Ser Lys Thr Gly Pro Gly Gln Lys Ala Ile Leu Phe Leu Pro
 140 145 150

ATG TCT GCT AAG AGC TGA TAA 471
 Met Ser Ala Lys Ser
 155

(17) INFORMATION FOR SEQ ID NO: 16

(i) SEQUENCE CHARACTERISTICS

- (A) LENGTH: 477 bases encoding 157 amino acids
- (B) TYPE: nucleic acid and amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE

- (A) DESCRIPTION: DNA encoding a protein

(v) FRAGMENT TYPE: entire sequence

(vi) IMMEDIATE SOURCE: constructed

(ix) FEATURE

- (D) INFORMATION: FGF-2 having surface loop residues 118 to 122 replaced with corresponding structural elements from bovine FGF-1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO 16:

ATG GCT GAA GGG GAA ATC ACC ACG CTG CCC GCC CTT CCG GAG GAT 45
 Met Ala Glu Gly Glu Ile Thr Thr Leu Pro Ala Leu Pro Glu Asp
 5 10 15

GGC GGC AGC GGC GCC TTC CCG CCC GGG CAC TTC AAG GAC CCC AAG 90
 Gly Gly Ser Gly Ala Phe Pro Pro Gly His Phe Lys Asp Pro Lys
 20 25 30

CGG CTG TAC TGC AAA AAC GGG GGC TTC TTC CTG CGC ATC CAC CCC 135
 Arg Leu Tyr Cys Lys Asn Gly Gly Phe Phe Leu Arg Ile His Pro
 35 40 45

- 56 -

GAC	GGC	CGA	GTT	GAC	GGG	GTC	CGG	GAG	AAG	AGC	GAC	CCT	CAC	ATC	180
Asp	Gly	Arg	Val	Asp	Gly	Val	Arg	Glu	Lys	Ser	Asp	Pro	His	Ile	
				50					55					60	
AAG	CTA	CAA	CTT	CAA	GCA	GAA	GAG	AGA	GGA	GTT	GTG	TCT	ATC	AAA	225
Lys	Leu	Gln	Leu	Gln	Ala	Glu	Glu	Arg	Gly	Val	Val	Ser	Ile	Lys	
				65					70					75	
GGA	GTG	TGT	GCT	AAC	CGG	TAC	CTG	GCT	ATG	AAG	GAA	GAT	GGA	AGA	270
Gly	Val	Cys	Ala	Asn	Arg	Tyr	Leu	Ala	Met	Lys	Glu	Asp	Gly	Arg	
				80					85					90	
TTA	CTG	GCT	TCT	AAA	TGT	GTT	ACG	GAT	GAG	TGT	TTC	TTT	TTT	GAA	315
Leu	Leu	Ala	Ser	Lys	Cys	Val	Thr	Asp	Glu	Cys	Phe	Phe	Phe	Glu	
				95					100					105	
CGA	TTG	GAA	TCT	AAT	AAC	TAC	AAT	ACT	TAC	CGG	TCT	AAA	AAG	CAT	360
Arg	Leu	Glu	Ser	Asn	Asn	Tyr	Asn	Thr	Tyr	Arg	Ser	Lys	Lys	His	
				110					115					120	
GCT	GAA	AAA	CAC	TGG	TAT	GTG	GCA	TTG	AAA	CGA	ACT	GGG	CAG	TAT	405
Ala	Glu	Lys	His	Trp	Tyr	Val	Ala	Leu	Lys	Arg	Thr	Gly	Gln	Tyr	
				125					130					135	
AAA	CTT	GGT	TCC	AAA	ACA	GGA	CCT	GGG	CAG	AAA	GCT	ATA	CTT	TTT	450
Lys	Leu	Gly	Ser	Lys	Thr	Gly	Pro	Gly	Gln	Lys	Ala	Ile	Leu	Phe	
				140					145					150	
CTT	CCA	ATG	TCT	GCT	AAG	AGC	TGA	TAA							477
Leu	Pro	Met	Ser	Ala	Lys	Ser									
				155											

BIBLIOGRAPHY

- Arakawa, T. and Fox G.M., Eur. Pat. Ap. Pub. No. 320,148 (1989).
- Baird, A., et al., *Proc. Nat. Acad. Sci. USA* 85: 2324-2328 (1988).
- Baird, A., and Böhlen, P., *Handbook of Exp. Pharmacol.* 95(1): 369-418, Springer, 1990.
- Bergonzoni, L., et al., Eur. Pat. Ap. Pub. No. 363,675 (1990).
- Davidson, J.M., et al., *J. Cell Bio.* 100: 1219-1227 (1985).
- Eriksson, E.A., et al., *Proc. Nat. Acad. Sci. USA* 88: 3441-3445 (1991).
- Feige, J.J. et al., *Proc. Nat. Acad. Sci. USA* 86: 3174-3178 (1989).
- Fiddes, J.C., et al., Eur. Pat. Ap. Pub. No. 298,723 (1989).
- Franco, W.P., U.S. Pat. No. 4,378,347, Mar. 29, 1983.
- Givol, D., and Yayon, A., *FASEB J.* 6: 3362-3369 (1992).
- Gospardarowicz, D., et al., *Proc. Nat. Acad. Sci.* 81: 6963-6967 (1984).
- Howden, G.F., and Silver, I.A., *Int. Endodontic J.* 13: 3-6 (1980).
- Hynes, T.R., et al., *Nature* 339: 73-76 (1989).
- Jaye, M., et al., *Biochim. Biophys. Acta* 1135: 185-199 (1992).
- Johnson, D.E., and Williams, L.T., *Adv. Can. Res.* 60: 1-41 (1993).
- Miyamoto, M., et al., *Mol. Cell. Biol.* 13: 4251-4259 (1993).
- Moscattelli, D., *J. Cell. Physiol.* 131: 123-130 (1987).
- Presta, M., et al., *B.B.R.C.* 185: 1098-1107 (1992).
- Seddon, A.P. et al., *Annals N.Y. Acad. Sci.* 638: 98-108 (1991).
- Seno, M., et al., Eur. Pat. Ap. Pub. Nos. 281,822' (1988) and 326,907 (1989).
- Seno, M., et al., *Eur. J. Biochem.* 188: 239-245 (1990).
- Tanaka, A., et al., *Proc. Natl. Acad. Sci. USA* 89: 8928-8932 (1992).
- Werner, S., et al., *Mol. Cell. Bio.* 12: 82-88 (1992).
- Yayon, A., et al., *EMBO J.* 11: 1885-1890 (1992).
- Zhang, J., et al., *Proc. Nat. Acad. Sci. USA* 88: 3446-3450 (1991).
- Zhu, H., et al., *Science* 251: 90-93 (1991).

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>26</u> , line <u>17</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 12301 Parklawn Drive Rockville, Maryland United States of America	
Date of deposit 17 September 1993	Accession Number A.T.C.C. 69417, 69418 69419, 69420
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
<div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black; margin: 0 10px;">For receiving Office use only</div> <div style="border: 1px solid black; padding: 5px; margin-top: 5px;"><input checked="" type="checkbox"/> This sheet was received with the international application</div> <div style="border: 1px solid black; padding: 5px; margin-top: 5px;">Authorized officer <u>SRU</u> Sigfried R. Hostad PCT International Division 703-305-3682</div>	<div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black; margin: 0 10px;">For International Bureau use only</div> <div style="border: 1px solid black; padding: 5px; margin-top: 5px;"><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div style="border: 1px solid black; padding: 5px; margin-top: 5px;">Authorized officer</div>

CLAIMS

1. A fibroblast growth factor structural analogue comprising a fibroblast growth factor having an amino acid sequence replacement in the sequence that corresponds to the surface loop that connects the ninth and tenth β -
5 strands in fibroblast growth factor-2, or the loop stems or anchor points, such that the overall secondary and tertiary structure of the protein is not significantly perturbed and the analogue binds to heparin, and wherein said analogue exhibits binding affinity to a native fibro-
10 blast growth factor receptor.
2. A structural analogue according to claim 1 wherein the binding specificity to a fibroblast growth factor receptor is different from the corresponding binding specificity of native factor from which the analogue is derived.
3. A fibroblast growth factor structural analogue according to claims 1 or 2 comprising a fibroblast growth factor having the amino acid sequence in the growth factor surface loop that connects the ninth to the tenth β -strand
5 replaced by an amino acid sequence that does not significantly perturb the overall folding of the molecule, so that the analogue binds to heparin and a member of the fibroblast growth factor receptor family.
4. A structural analogue according to claim 3 comprising a fibroblast growth factor-2 structural analogue.
5. A structural analogue according to claim 3 wherein the amino acid sequence in the growth factor surface loop is replaced by a corresponding amino acid sequence derived from another fibroblast growth factor surface loop or an
5 interleukin surface loop.

6. A structural analogue according to claim 5 wherein the binding specificity of the analogue exhibits similarities to the fibroblast growth factor from which the loop is derived.

7. A structural analogue according to claims 5 or 6 wherein a corresponding amino acid surface loop sequence from another fibroblast growth factor replaces sequences selected from the group consisting of:

- (a) fibroblast growth factor-1 sequence Lys115-Lys-
5 116-His117-Ala118-Glu119-Lys120-Asn121;
- (b) fibroblast growth factor-2 sequence Arg118-Lys119-Tyr120-Thr121-Ser122;
- (c) fibroblast growth factor-3 sequence Arg-Leu-Tyr-Arg-Thr-Val-Ser-Ser-Thr-Pro-Gly-Ala-Arg-Arg-Gln-Pro-
10 Ser-Ala-Glu-Arg-Leu;
- (d) fibroblast growth factor-4 sequence Tyr-Lys-Tyr-Pro-Gly;
- (e) fibroblast growth factor-5 sequence Ala-Ile-His-Arg-Thr-Glu-Lys-Thr-Gly-Arg-Glu;
- 15 (f) fibroblast growth factor-6 sequence Asp-Leu-Tyr-Gln-Gly;
- (g) fibroblast growth factor-7 sequence Ala-Lys-Trp-Thr-His-Asn-Gly-Gly-Glu;
- (h) fibroblast growth factor-8 sequence Ala-Lys-
20 Tyr-Glu-Gly; and
- (i) fibroblast growth factor-9 sequence Asn-Leu-Tyr-Lys-His-Val-Asp-Thr-Gly-Arg-Arg.

8. A structural analogue according to claim 7 wherein the corresponding amino acid sequence is Lys115-Lys116-His117-Ala118-Glu119-Lys120-His121 from bovine fibroblast growth factor-1, Ala231-Gln232-Phe233-Pro234-Asn235 from human interleukin-1 β , or Ala154-Lys155-Trp156-Thr157-His158-Asn159-Gly160-Gly161-Glu162 from fibroblast growth factor-


9. A structural analogue according to claim 8 wherein said amino acid sequence replacement occurs between residues 112 and 128.

10. A fibroblast growth factor from the family of fibroblast growth factor-1 or fibroblast growth factors 3 to 9 wherein the sequences corresponding to the loop in fibroblast growth factor-2 connecting the ninth and tenth β -strands in fibroblast growth factor-2 is replaced by a
5 corresponding loop from another fibroblast growth factor or an interleukin 1.

FGF-2 Asn Asn Tyr Asn Thr Tyr Arg Ser Arg Lys Tyr Thr Ser Trp Tyr Val Ala Leu Lys
 110 115 120 125
 (Ser)
 FGF-1 Asn His Tyr Asn Thr Tyr Ile Ser Lys Lys His Ala Glu Lys Asn Trp Phe Val Gly Leu Lys
 (His)
 IL-18 Asn Asn Lys Leu Glu Phe Glu Ser Ala Gln Phe Pro Asn Trp Tyr Ile Ser Thr Ser

Figure 1

Figure 2




**Figure 3**

Figure 4






Figure 5

Figure 6





Figure 7

**Figure 8**




Figure 9A




Figure 9B

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/10800

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 15/18, 15/19, 15/24, 15/25; A61K 38/18, 38/19, 38/20

US CL : 435/69.4, 69.5, 69.52, 69.7; 530/350, 351, 399

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.4, 69.5, 69.52, 69.7; 530/350, 351, 399

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS and DIALOG (files 5, 155, 351, 357, 358) search terms: FGF, fibroblast growth factor, IL-1, interleukin, surface loop, beta strand

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EMBO JOURNAL, Volume 5, No. 10, issued 1986, J.A. Abraham et al, "Human basic fibroblast growth factor: nucleotide sequence and genomic organization", pages 2523-2528, especially figures 2 and 3.	1-10
X	EUROPEAN JOURNAL OF BIOCHEMISTRY, Volume 188, issued 1990, M. Seno et al, "Carboxyl-terminal structure of basic fibroblast growth factor significantly contributes to its affinity for heparin", pages 239-245, see entire document.	1,3



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:		T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A	document defining the general state of the art which is not considered to be of particular relevance	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E	earlier document published on or after the international filing date	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	A	document member of the same patent family
O	document referring to an oral disclosure, use, exhibition or other means		
P	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

01 DECEMBER 1994

Date of mailing of the international search report

11 JAN 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MARIANNE PORTA ALLEN

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/10800

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, Volume 185, No. 3, issued 30 June 1992, M. Presta et al, "Structure-function relationship of basic fibroblast growth factor: site-directed mutagenesis of a putative heparin-binding and receptor-binding region", pages 1098-1107, see entire document.	1-4, 9
X	BIOCHEMISTRY, Volume 30, issued 1991, W.F. Heath et al, "Mutations in the heparin-binding domains of human basic fibroblast growth factor alter its biological activity", pages 5608-5615, see entire document.	1-6, 9